

Coincidence of topological Jacobson radicals in topological algebras

MART ABEL, MATI ABEL, AND PAUL TAMMO

ABSTRACT. Several classes of topological algebras for which the left topological Jacobson radical coincides with the right topological Jacobson radical are described.

1. Introduction

The notion of a *topological (Jacobson) radical* was introduced in [3] as the intersection of kernels of all continuous irreducible representations of a topological algebra A on all Hausdorff topological linear spaces X in case when the module multiplication over A on X is commutative (that is, when $ax = xa$ for all $a \in A$ and $x \in X$). In general, two different topological Jacobson radicals exist: the left topological Jacobson radical defined by the continuous irreducible representations of A on X , and the right topological Jacobson radical defined by the continuous irreducible antirepresentations of A on X . A class of topological algebras, for which the left and right topological radicals coincide, has been described in [1]. Several new classes of topological algebras, for which the left topological Jacobson radical coincides with the right topological Jacobson radical, are given in the present paper.

Let \mathbb{K} denote one of the fields \mathbb{R} or \mathbb{C} of real or complex numbers, respectively. An associative algebra A over \mathbb{K} is a *topological algebra over \mathbb{K}* (shortly, a *topological algebra*) if A is a topological linear space over \mathbb{K} (with respect to the same linear operations) and the multiplication in A is separately continuous. We denote by θ_A the zero element of A .

By an ideal we always mean a proper ideal. A left (right or two-sided) ideal I of A is *regular* (sometimes also called *modular*) if there exists an element $u \in A$ such that $a - au \in I$ ($a - ua \in I$ or $a - au, a - ua \in I$, respectively) for

Received June 30, 2016.

2010 *Mathematics Subject Classification.* 46H05, 46H10.

Key words and phrases. Topological algebras, topological Jacobson radicals.

<http://dx.doi.org/10.12697/ACUTM.2017.21.16>

all $a \in A$. Such an element u is called a *right (left or two-sided, respectively) regular unit for I* . It is known that $u \notin I$ (see [2]).

If M is a maximal regular left (right) ideal of A with a right (left, respectively) regular unit u , then the two-sided ideal $P = \{p \in A: pA \subset M\}$ ($P = \{p \in A: Ap \subset M\}$, respectively) is called the *primitive ideal defined by M* . If M is closed, then P is closed (see [2]).

A topological algebra A is called *left (right) simplicial* if every closed regular left (right, respectively) ideal of A is contained in a closed maximal regular left (right, respectively) ideal of A , and *simplicial* if it is both left and right simplicial.

Let A be a topological algebra (or a topological ring in general). An element r of A is *left quasi-invertible* if there is an element $p \in A$ such that $p \circ r = p + r - pr = \theta_A$. An element r of A is *left topologically quasi-invertible* if there is a net $(p_\lambda)_{\lambda \in \Lambda}$ in A such that the net $(p_\lambda \circ r)_{\lambda \in \Lambda}$ converges to θ_A . We denote by $\text{Qinv}_l A$ the set of all left quasi-invertible elements of A and by $\text{Tqinv}_l A$ the set of all left topologically quasi-invertible elements of A . The sets of all right quasi-invertible elements $\text{Qinv}_r A$ of A and all right topologically quasi-invertible elements $\text{Tqinv}_r A$ of A are defined similarly. We say that a topological algebra is *advertive* if $\text{Tqinv}_r A = \text{Qinv}_r A$ and $\text{Tqinv}_l A = \text{Qinv}_l A$.

Let A be an algebra over \mathbb{K} and M a linear space over \mathbb{K} . The space M is called a *left (right) A -module* if the module multiplication $(a, m) \rightarrow am$ of $A \times M$ into M ($(m, a) \rightarrow ma$ of $M \times A$ into M , respectively) is a bilinear map which satisfies the condition $a_1(a_2m) = (a_1a_2)m$ ($(ma_1)a_2 = m(a_1a_2)$, respectively) for each $a_1, a_2 \in A$ and $m \in M$. If M' is a subspace of a left (right) A -module M and for any $m \in M'$ and $a \in A$ the product am (ma , respectively) is in M' , then M' is a *left (right, respectively) A -submodule* of M . In particular, when A is a topological algebra over \mathbb{K} (with separately continuous multiplication) and M is a topological linear space over \mathbb{K} in which the module multiplication is separately continuous, M is called a *topological left (right) A -module*. A left (right) A -module X is *nontrivial* if $AX \neq \{\theta_X\}$ ($XA \neq \{\theta_X\}$, respectively) and X is *irreducible* if it is nontrivial and the only A -submodules of X are X and $\{\theta_X\}$.

Let X be a topological linear space over \mathbb{K} , and let $\mathcal{L}(X)$ be the set of all continuous linear maps on X . If we define the addition of elements and the multiplication over \mathbb{K} in $\mathcal{L}(X)$ point-wise and the multiplication of elements in $\mathcal{L}(X)$ by the composition, then $\mathcal{L}(X)$ is an algebra (non-commutative in general) over \mathbb{K} .

Remind that an *antihomomorphism* is a map $\pi' : A \rightarrow B$ between two algebras that preserves addition and multiplication by scalars but reverses the order of multiplication, that is, $\pi'(xy) = \pi'(y)\pi'(x)$ for all $x, y \in A$. Any homomorphism π (antihomomorphism π') of an algebra A into $\mathcal{L}(X)$ is called a *representation (antirepresentation, respectively) of A on X* .

Every representation π (antirepresentation π') of A on X defines on X a left (right, respectively) module multiplication \cdot_π ($\cdot_{\pi'}$, respectively) if we put $a \cdot_\pi x = \pi(a)(x)$ ($x \cdot_{\pi'} a = \pi'(a)(x)$, respectively) for each $a \in A$ and $x \in X$. In this case X becomes a left (right, respectively) A -module. We denote it by X_π ($X_{\pi'}$, respectively). A representation π (antirepresentation π' , respectively) of A on X is *irreducible* if X_π ($X_{\pi'}$, respectively) is an irreducible left (right, respectively) A -module. Let $x_0 \in X$ and define $\varrho_{\pi, x_0} : A \rightarrow X$ by

$$\varrho_{\pi, x_0}(a) = a \cdot_\pi x_0$$

if X is a left A -module and $\varrho_{\pi', x_0} : A \rightarrow X$ by

$$\varrho_{\pi', x_0}(a) = x_0 \cdot_{\pi'} a$$

if X is a right A -module, for all $a \in A$. If X is a left A -module and, thus, π is a representation of A on X , then

$$\ker \varrho_{\pi, x_0} = \{a \in A : a \cdot_\pi x_0 = \theta_A\}$$

is a left ideal of A . If X is a right A -module and, thus, π' is an antirepresentation of A on X , then

$$\ker \varrho_{\pi', x_0} = \{a \in A : x_0 \cdot_{\pi'} a = \theta_A\}$$

is a right ideal of A . We say that x_0 is a *left (right) cyclic element* if $A \cdot_\pi x_0 = X$ ($x_0 \cdot_{\pi'} A = X$, respectively). Moreover, denote by

$$\text{id}_r(x_0) = \{e \in A : e \cdot_\pi x_0 = x_0\}$$

and

$$\text{id}_l(x_0) = \{e \in A : x_0 \cdot_{\pi'} e = x_0\}$$

the sets of one-sided units for an element x_0 .

The intersection $\text{Rad}(A)$ of kernels of all irreducible representations of an algebra A on linear spaces is called the *Jacobson radical of A* . If there are no irreducible representations of A , then A is called a *radical algebra* (in this case $\text{Rad}(A) = A$) and if $\text{Rad}(A) = \{\theta_A\}$, then A is called a *semi-simple algebra*.

We endow $\mathcal{L}(X)$ with the topology τ of simple convergence. A base of neighbourhoods of zero of τ consists of sets

$$T(S, O) = \{L \in \mathcal{L}(X) : L(S) \subset O\},$$

where S is a finite subset of X and O is a neighbourhood of zero in X . If X is a Hausdorff linear space, then $\mathcal{L}(X)$ is a Hausdorff algebra with separately continuous multiplication. We denote this algebra by $\mathcal{L}_\tau(X)$.

Let A be a topological algebra, M a closed maximal regular left (right) ideal of A , A/M the quotient space of A modulo M , and κ the canonical homomorphism from A onto A/M . It is easy to see that A/M is a Hausdorff linear space in the quotient topology if we define the addition and the multiplication over \mathbb{K} in A/M as usual, and A/M is a topological left (right,

respectively) A -module if we define the module multiplication in A/M over A by $ax = \kappa(aa')$ for each $a \in A$ and $x \in A/M$, where a' is an arbitrary element of the M -coset of x .

Let again A be a topological algebra. For each $a \in A$ and each closed maximal regular left (right) ideal M of A , let L_a^M (R_a^M , respectively) be the map from A/M into A/M defined by $L_a^M(x) = ax$ ($R_a^M(x) = xa$, respectively) for each $x \in A/M$, and let L_M (R_M , respectively) be the map from A into $\mathcal{L}(A/M)$ defined by $L_M(a) = L_a^M$ ($R_M(a) = R_a^M$, respectively) for each $a \in A$. Then L_M (R_M , respectively) is a representation (antirepresentation, respectively) of A on A/M , $a \cdot_{L_M} x = ax$ ($a \cdot_{R_M} x = xa$, respectively) for each $a \in A$ and $x \in A/M$, and the left (right, respectively) A -module $(A/M)_{L_M}$ ($(A/M)_{R_M}$, respectively) is irreducible (see [7], Proposition 5, page 120). Hence, L_M (R_M , respectively) is an irreducible representation (antirepresentation, respectively) of A on A/M . Thus

$$\ker L_M = \{a \in A : aA \subset M\} \quad (\ker R_M = \{a \in A : Aa \subset M\})$$

is a primitive ideal. More precisely, we call the set $\ker L_M$ ($\ker R_M$, respectively) the *primitive ideal defined by the maximal regular left (right, respectively) ideal M of A* . To show that L_M is continuous, let U be an arbitrary neighbourhood of zero in $\mathcal{L}_\tau(A/M)$. Then there exists a neighbourhood O of zero of A/M and a finite subset S of A/M such that

$$T = \{L \in \mathcal{L}_\tau(A/M) : L(S) \subset O\} \subset U.$$

Since the multiplication in A/M over A is separately continuous, then for each $s \in S$ there exists a neighbourhood V_s of θ_A such that $V_s s \subset O$ ($sV_s \subset O$). Let now

$$V = \bigcap_{s \in S} V_s.$$

Then $VS \subset O$ ($SV \subset O$) and V is a neighbourhood of zero in A . Since $L_M(v)(S) = L_v^M(S) = vS \subset O$ for each $v \in V$, one has $L_M(V) \subset T \subset U$. Therefore, L_M is a continuous map (because L_M is linear). Hence, L_M is a continuous irreducible representation of A on A/M and $\ker L_M$ is a closed two-sided ideal of A . Similarly, we can show that the antirepresentation R_M is continuous.

As defined in [1], the *left (right) topological radical* of a topological algebra A is the intersection $\text{rad}_l(A)$ ($\text{rad}_r(A)$) of kernels of all continuous irreducible representations (antirepresentations) of A on all Hausdorff linear spaces. If A has no continuous irreducible representations (antirepresentations) at all, then we say that A is a *left (right, respectively) topologically radical algebra* (in this case $\text{rad}_l(A) = \text{rad}_r(A) = A$). If $\text{rad}_l(A) \neq A$ ($\text{rad}_r(A) \neq A$), then we say that A is a *left (right) topologically nonradical algebra*, and if $\text{rad}_l(A) = \{\theta_A\}$ ($\text{rad}_r(A) = \{\theta_A\}$), then A is a *left (right, respectively) topologically semi-simple algebra*.

2. Description of one-sided topological radicals

The left-sided case of the following lemma is Proposition 4 in [7], page 120. Here we adopt the notation and give for completeness the proof for the right-sided case.

Lemma 1. *Let X be a left (right) A -module, let π (π' , respectively) be a representation (antirepresentation, respectively) of A on X , and let $x_0 \in X \setminus \{\theta_X\}$.*

- a) *Each element of $\text{id}_r(x_0)$ ($\text{id}_l(x_0)$) is a right (left, respectively) regular unit for the left ideal $\ker \varrho_{\pi, x_0}$ (right ideal $\ker \varrho_{\pi', x_0}$, respectively).*
- b) *If x_0 is a left (right) cyclic element, then $\text{id}_r(x_0)$ ($\text{id}_l(x_0)$) is non-void and $\ker \varrho_{\pi, x_0}$ is a regular left (ker ϱ_{π', x_0} is a regular right, respectively) ideal of A .*
- c) *If X is irreducible, then x_0 is a left (right, respectively) cyclic element and $\ker \varrho_{\pi, x_0}$ (ker ϱ_{π', x_0} , respectively) is a maximal regular left (right, respectively) ideal.*

Proof. a) Let $e \in \text{id}_l(x_0)$. Then, for all $a \in A$,

$$x_0 \cdot_{\pi'} (a - ea) = x_0 \cdot_{\pi'} a - x_0 \cdot_{\pi'} a = \theta_A$$

and so $a - ea \in \ker \varrho_{\pi', x_0}$.

b) Since x_0 is a right cyclic element, $x_0 \cdot_{\pi'} A = X$. So there is an element $e \in A$ such that $x_0 \cdot_{\pi'} e = x_0$, that is, $e \in \text{id}_l(x_0)$. By the statement a), e is a left regular unit for $\ker \varrho_{\pi', x_0}$ and therefore $\ker \varrho_{\pi', x_0}$ is a regular right ideal of A .

c) Let X be irreducible. Since $x_0 \cdot_{\pi'} A$ is an A -submodule, either $x_0 \cdot_{\pi'} A = \{\theta_A\}$ or $x_0 \cdot_{\pi'} A = A$. If $x_0 \cdot_{\pi'} A = \{\theta_A\}$, then $N = \{x \in X : x \cdot_{\pi'} A = \{\theta_A\}\}$ is a non-zero A -submodule, and so $N = X$ because X is irreducible. In this case $X \cdot_{\pi'} A = \{\theta_A\}$, but this is impossible since X is non-trivial. Consequently, $x_0 \cdot_{\pi'} A = X$ and x_0 is right cyclic. By the statement b), $\ker \varrho_{\pi', x_0}$ is a regular right ideal. Let J be a right ideal of A with $\ker \varrho_{\pi', x_0} \subsetneq J$. Now $x_0 \cdot_{\pi'} J$ is an A -submodule and for every $j \in J \setminus \ker \varrho_{\pi', x_0}$ we have $x_0 \cdot_{\pi'} j \neq \theta_A$ by the definition of $\ker \varrho_{\pi', x_0}$, which means that $x_0 \cdot_{\pi'} J \neq \{\theta_A\}$. Therefore $x_0 \cdot_{\pi'} J = X$. So there exists an element $e \in J \cap \text{id}_l(x_0)$. By the statement a), e is a left regular unit for $\ker \varrho_{\pi', x_0}$ and, thus, also for J . But then $J = A$ since $e \in J$. The proof for the left-sided case is similar. \square

Theorem 1 in [4] considers the case when the A -module X is commutative. Here we examine the case when the multiplication over A is not necessarily commutative.

We denote by $m_l(A)$ ($m_r(A)$ or $m_t(A)$) the set of all closed maximal regular left (right or two-sided, respectively) ideals of A . In addition, $\pi_l(A)$

$(\pi_r(A))$ is the set of all closed primitive ideals of A defined by closed maximal regular left (right, respectively) ideals.

Theorem 2. *Let A be a right (left) topologically nonradical algebra. Then $\text{rad}_r(A)$ ($\text{rad}_l(A)$, respectively) is equal to*

- a) *the intersection of all closed maximal regular right (left, respectively) ideals of A ;*
- b) *the intersection of all primitive ideals of A which are defined by closed maximal regular right (left, respectively) ideals of A .*

That is,

$$\text{rad}_k(A) = \bigcap_{P \in \pi_k(A)} P = \bigcap_{M \in m_k(A)} M$$

where $k = r$ ($k = l$, respectively).

Hence, $\text{rad}_r(A)$ ($\text{rad}_l(A)$, respectively) is closed in A .

Proof. Since $\ker R_M \subset M$ for each $M \in m_r(A)$, we have

$$\text{rad}_r(A) \subset \bigcap_{M \in m_r(A)} \ker R_M \subset \bigcap_{M \in m_r(A)} M.$$

To show the converse inclusion, let $a \in \cap\{M : M \in m_r(A)\}$. If $a \notin \text{rad}_r(A)$, then there is a continuous irreducible antirepresentation π' of A on some Hausdorff linear space X such that $a \notin \ker \pi'$. Therefore there exists an element $x_0 \in X$ such that $\pi'(a)(x_0) \neq \theta_X$. Hence $a \notin I = \ker \varrho_{\pi',x_0}$. By Lemma 1, $\ker \varrho_{\pi',x_0}$ is a maximal regular right ideal of A . To show that $\ker \varrho_{\pi',x_0}$ is closed in A , we show that the map ϱ_{π',x_0} is continuous on A . To this end, let a_0 be an arbitrary element in A , let O be a neighbourhood of $\pi'(a_0)(x_0)$ in X , let O_X be a neighbourhood of zero in X such that $O = \pi'(a_0)(x_0) + O_X$, and let

$$O(\pi'(a_0)) = \{L \in \mathcal{L}_\tau(X) : (L - \pi'(a_0))(x_0) \in O_X\}.$$

Since $O(\pi'(a_0))$ is a neighbourhood of $\pi'(a_0)$ in $\mathcal{L}_\tau(X)$ and π' is a continuous antirepresentation of A on X , there is a neighbourhood $O(a_0)$ of a_0 in A such that $\pi'(O(a_0)) \subset O(\pi'(a_0))$. If now $b \in O(a_0)$, then

$$\varrho_{\pi',x_0}(b) = \pi'(b)(x_0) \in \pi'(a_0)(x_0) + O_X = O.$$

Therefore ϱ_{π',x_0} is continuous. It means that I is a closed maximal regular right ideal of A which does not contain a , but this is not possible. Consequently, $a \in \text{rad}_r(A)$. Hence the statement a) holds. Since $\ker R_M$ is a closed primitive ideal of A for each $M \in m_r(A)$, the statement b) holds too.

The proof for left topological radicals is similar. □

Note that $\text{Rad}(A) \subset \text{rad}_l(A) \cap \text{rad}_r(A)$.

3. Coincidence of left and right topological radicals

The description of the class of unital topological algebras in which all maximal one-sided ideals are closed has been given in [5], and the class in which all maximal two-sided ideals are closed is described in [6] (for the nonunital case, see [3]). In these cases $\text{rad}_l(A) = \text{rad}_r(A)$ by Theorem 2. This is the case when A is, e.g., a Q -algebra (that is, the set of all left quasi-invertible elements and the set of all right quasi-invertible elements of A are open). Here we consider the case when maximal regular ideals are not necessarily closed. A class of topological algebras for which one-sided topological radicals coincide has been described in [1].

Theorem 3. *Let A be a topological algebra and $\kappa_P: A \rightarrow A/P$ be the quotient map for every $P \in \pi(A)$. Then $\text{rad}_l(A) = \text{rad}_r(A)$ if and only if*

$$\bigcap_{P \in \pi_l(A)} \kappa_P^{-1}(\text{rad}_l(A/P)) = \bigcap_{Q \in \pi_r(A)} \kappa_Q^{-1}(\text{rad}_r(A/Q)). \tag{1}$$

Proof. Note that

$$\begin{aligned} \text{rad}_l(A) &= \bigcap_{M \in m_l(A)} M = \bigcap_{P \in \pi_l(A)} \bigcap_{M \in m_l(A/P)} \kappa_P^{-1}(M) \\ &= \bigcap_{P \in \pi_l(A)} \kappa_P^{-1} \left(\bigcap_{M \in m_l(A/P)} M \right) \\ &= \bigcap_{P \in \pi_l(A)} \kappa_P^{-1}(\text{rad}_l(A/P)) \end{aligned}$$

by Theorem 2 and Theorem 12 in [2]. Similarly,

$$\text{rad}_r(A) = \bigcap_{Q \in \pi_r(A)} \kappa_Q^{-1}(\text{rad}_r(A/Q))$$

and, therefore, $\text{rad}_l(A) = \text{rad}_r(A)$ if and only if (1) holds. □

Corollary 4. *Let A be a topological algebra. If $\pi_l(A) = \pi_r(A)$ and $\text{rad}_l(A/P) = \text{rad}_r(A/P)$ for every $P \in \pi_l(A)$, then $\text{rad}_l(A) = \text{rad}_r(A)$.*

Proof. If $\pi_l(A) = \pi_r(A)$ and $\text{rad}_l(A/P) = \text{rad}_r(A/P)$ for every $P \in \pi_l(A)$, then (1) holds and so, by Theorem 3, $\text{rad}_l(A) = \text{rad}_r(A)$. □

Theorem 5. *Let A be a simplicial topological algebra and I a closed two-sided ideal of A . Then A/I is simplicial.*

Proof. Let \mathcal{J} be a closed left (right) ideal of A/I . Then $\kappa_I^{-1}(\mathcal{J})$ is a closed left (respectively, right) ideal of A (by Corollary 2 in [2]). Since A is simplicial, there is $M \in m_l(A)$ ($M \in m_r(A)$, respectively) such that $\kappa_I^{-1}(\mathcal{J}) \subset M$. Therefore, $\mathcal{J} = \kappa_I(\kappa_I^{-1}(\mathcal{J})) \subset \kappa_I(M)$. Since $I \subset M$, then $\kappa_I(M)$ is a closed maximal regular left (right, respectively) ideal of A/I , by the statement a)

of Corollary 2 in [2]. Hence, every closed left (right, respectively) ideal of A/I is contained in a closed maximal regular left (right, respectively) ideal of A/I . \square

If all closed maximal regular one-sided ideals in a topological algebra A are two-sided ideals, then $\pi_l(A) = \pi_r(A)$.

Corollary 6. *Let A be a topological algebra. If $\pi_l(A) = \pi_r(A)$ and A/P is simplicial and advertive for every $P \in \pi_l(A)$, then $\text{rad}_l(A) = \text{rad}_r(A)$.*

Proof. By the assumption, A/P is simplicial and advertive for every $P \in \pi_l(A)$, and therefore, by Theorem 1 in [1], $\text{rad}_l(A/P) = \text{rad}_r(A/P)$ for every $P \in \pi_l(A)$. Since also $\pi_l(A) = \pi_r(A)$, by Corollary 4, $\text{rad}_l(A) = \text{rad}_r(A)$. \square

Corollary 7. *Let A be a simplicial topological algebra. If $\pi_l(A) = \pi_r(A)$ and A/P is advertive for every $P \in \pi_l(A)$, then $\text{rad}_l(A) = \text{rad}_r(A)$.*

Proof. By Theorem 5, A/P is simplicial for every $P \in \pi_l(A)$, since A is simplicial. Now, by Corollary 6, $\text{rad}_l(A) = \text{rad}_r(A)$. \square

Corollary 8. *Let A be a simplicial topological algebra. If $\pi_l(A) = \pi_r(A)$ and for every $P \in \pi_l(A)$ there exists an advertive subalgebra $B_P \subset A$ such that $A = P \oplus B_P$, then $\text{rad}_l(A) = \text{rad}_r(A)$.*

Proof. Let A be a simplicial topological algebra. If $A = P \oplus B_P$ for every $P \in \pi_l(A)$, then $A/P \cong B_P$ for every $P \in \pi_l(A)$ and, by Corollary 7, $\text{rad}_l(A) = \text{rad}_r(A)$. \square

Acknowledgement

The research is partially supported by institutional research funding IUT20-57 of the Estonian Ministry of Education and Research.

References

- [1] Mart Abel and Mati Abel, *On a problem of Bertram Yood*, Topol. Algebra Appl. **2**(1) (2014), 1–4.
- [2] Mart Abel, Mati Abel, and Paul Tammo, *Descriptions of all closed maximal one-sided ideals in topological algebras*, Proc. ICTAA 2015, accepted.
- [3] Mati Abel, *Description of topological algebras without a unit in which all maximal regular two-sided ideals are closed*, Rev. Bull. Calcutta Math. Soc. **20** (2012), 1–10.
- [4] Mati Abel, *Descriptions of the topological radical in topological algebras*, in: General Topological Algebras (Tartu, 1999), Math. Stud. (Tartu) **1**, Est. Math. Soc., Tartu, 2001, pp. 25–31.
- [5] Mati Abel, *Unital topological algebras in which all maximal one-sided ideals are closed*, in: International Conference on Topological Algebras and their Applications (ICTAA, 2013), Math. Stud. (Tartu) **6**, Est. Math. Soc., Tartu, 2014, pp. 49–54.

- [6] Mati Abel and K. Jarosz, *Topological algebras in which all maximal two-sided ideals are closed*, in: Topological Algebras, their Applications, and Related Topics, Banach Center Publ. **67**, Polish Acad. Sci., Warsaw, 2005, pp. 35–43.
- [7] F. F. Bonsall and J. Duncan, *Complete Normed Algebras*, Ergebnisse der Mathematik und ihrer Grenzgebiete **80**. Springer-Verlag, New York–Heidelberg, 1973.

SCHOOL OF DIGITAL TECHNOLOGIES, TALLINN UNIVERSITY, 29 NARVA STR., 10120 TALLINN, ESTONIA; INSTITUTE OF MATHEMATICS AND STATISTICS, UNIVERSITY OF TARTU, 2 J. LIIVI STR., 50409 TARTU, ESTONIA
E-mail address: `mart.abel@tlu.ee`, `mart.abel@ut.ee`

INSTITUTE OF MATHEMATICS AND STATISTICS, UNIVERSITY OF TARTU, 2 J. LIIVI STR., 50409 TARTU, ESTONIA
E-mail address: `mati.abel@ut.ee`

INSTITUTE OF MATHEMATICS AND STATISTICS, UNIVERSITY OF TARTU, 2 J. LIIVI STR., 50409 TARTU, ESTONIA
E-mail address: `paul.tammo@ut.ee`